

Higher-order generalized Cahn–Hilliard equations

Laurence Cherfils¹, Alain Miranville², Shuiran Peng² and
Wen Zhang ³

¹Université de La Rochelle, Laboratoire Mathématiques, Image et Applications
Avenue Michel Crépeau, F-17042 La Rochelle Cedex, France

²Université de Poitiers, Laboratoire de Mathématiques et Applications, UMR CNRS 7348 - SP2MI
Boulevard Marie et Pierre Curie - Téléport 2, F-86962 Chasseneuil Futuroscope Cedex, France

³University of Xiamen, School of Mathematical Sciences, 361005 Xiamen, Fujian, China

Received 23 September 2016, appeared 31 January 2017

Communicated by Dimitri Mugnai

Abstract. Our aim in this paper is to study higher-order (in space) anisotropic generalized Cahn–Hilliard models. In particular, we obtain well-posedness results, as well as the existence of the global attractor. Such models can have applications in biology, image processing, etc. We also give numerical simulations which illustrate the effects of the higher-order terms on the anisotropy.

Keywords: generalized Cahn–Hilliard equation, higher-order models, anisotropy, well-posedness, global attractor, numerical simulations.

2010 Mathematics Subject Classification: 35K55, 35J60.

1 Introduction

The Cahn–Hilliard equation,

$$\frac{\partial u}{\partial t} + \Delta^2 u - \Delta f(u) = 0, \quad (1.1)$$

plays an essential role in materials science and describes important qualitative features of two-phase systems related with phase separation processes, assuming isotropy and a constant temperature. This can be observed, e.g., when a binary alloy is cooled down sufficiently. One then observes a partial nucleation (i.e., the apparition of nuclides in the material) or a total nucleation, the so-called spinodal decomposition: the material quickly becomes inhomogeneous, forming a fine-grained structure in which each of the two components appears more or less alternatively. In a second stage, which is called coarsening, occurs at a slower time scale and is less understood, these microstructures coarsen. Such phenomena play an essential role in the mechanical properties of the material, e.g., strength. We refer the reader to, e.g., [8, 9, 16, 20, 29, 30, 32, 33, 38, 39] for more details.

 Corresponding author. Email: zhangwenmath@126.com

Here, u is the order parameter (e.g., a density of atoms) and f is the derivative of a double-well potential F . A thermodynamically relevant potential F is the following logarithmic function which follows from a mean-field model:

$$F(s) = \frac{\theta_c}{2}(1-s^2) + \frac{\theta}{2} \left[(1-s) \ln \left(\frac{1-s}{2} \right) + (1+s) \ln \left(\frac{1+s}{2} \right) \right], \quad s \in (-1, 1), \quad 0 < \theta < \theta_c, \quad (1.2)$$

i.e.,

$$f(s) = -\theta_c s + \frac{\theta}{2} \ln \frac{1+s}{1-s}, \quad (1.3)$$

although such a function is very often approximated by regular ones, typically,

$$F(s) = \frac{1}{4}(s^2 - 1)^2, \quad (1.4)$$

i.e.,

$$f(s) = s^3 - s. \quad (1.5)$$

Now, it is interesting to note that the Cahn–Hilliard equation and some of its variants are also relevant in other phenomena than phase separation. We can mention, for instance, population dynamics (see [18]), tumor growth (see [4] and [26]), bacterial films (see [27]), thin films (see [41] and [44]), image processing (see [5, 6, 10, 12, 19]) and even the rings of Saturn (see [45]) and the clustering of mussels (see [31]).

In particular, several such phenomena can be modeled by the following generalized Cahn–Hilliard equation:

$$\frac{\partial u}{\partial t} + \Delta^2 u - \Delta f(u) + g(x, u) = 0. \quad (1.6)$$

We studied in [35] and [36] (see also [4, 12, 17, 21]) this equation.

The Cahn–Hilliard equation is based on the so-called Ginzburg–Landau free energy,

$$\Psi_{\text{GL}} = \int_{\Omega} \left(\frac{1}{2} |\nabla u|^2 + F(u) \right) dx, \quad (1.7)$$

where Ω is the domain occupied by the system (we assume here that it is a bounded and regular domain of \mathbb{R}^n , $n = 1, 2$ or 3 , with boundary Γ). In particular, in (1.7), the term $|\nabla u|^2$ models short-ranged interactions. It is however interesting to note that such a term is obtained by truncation of higher-order ones (see [9]); it can also be seen as a first-order approximation of a nonlocal term accounting for long-ranged interactions (see [22] and [23]).

G. Caginalp and E. Esenturk recently proposed in [7] (see also [11]) higher-order phase-field models in order to account for anisotropic interfaces (see also [28, 42, 47] for other approaches which, however, do not provide an explicit way to compute the anisotropy). More precisely, these authors proposed the following modified free energy, in which we omit the temperature:

$$\Psi_{\text{HOGL}} = \int_{\Omega} \left(\frac{1}{2} \sum_{i=1}^k \sum_{|\alpha|=i} a_{\alpha} |\mathcal{D}^{\alpha} u|^2 + F(u) \right) dx, \quad k \in \mathbb{N}, \quad (1.8)$$

where, for $\alpha = (k_1, \dots, k_n) \in (\mathbb{N} \cup \{0\})^n$,

$$|\alpha| = k_1 + \dots + k_n$$

and, for $\alpha \neq (0, \dots, 0)$,

$$\mathcal{D}^{\alpha} = \frac{\partial^{|\alpha|}}{\partial x_1^{k_1} \dots \partial x_n^{k_n}}$$

(we agree that $\mathcal{D}^{(0,\dots,0)}v = v$). The corresponding higher-order Cahn–Hilliard equation then reads

$$\frac{\partial u}{\partial t} - \Delta \sum_{i=1}^k (-1)^i \sum_{|\alpha|=i} a_\alpha \mathcal{D}^{2\alpha} u - \Delta f(u) = 0. \quad (1.9)$$

We studied in [13] and [14] the corresponding isotropic model which reads

$$\frac{\partial u}{\partial t} - \Delta P(-\Delta)u - \Delta f(u) = 0, \quad (1.10)$$

where

$$P(s) = \sum_{i=1}^k a_i s^i, \quad a_k > 0, \quad k \in \mathbb{N}, \quad s \in \mathbb{R}.$$

The anisotropic model (1.9) is treated in [15].

Our aim in this paper is to study the higher-order generalized Cahn–Hilliard model

$$\frac{\partial u}{\partial t} - \Delta \sum_{i=1}^k (-1)^i \sum_{|\alpha|=i} a_\alpha \mathcal{D}^{2\alpha} u - \Delta f(u) + g(x, u) = 0. \quad (1.11)$$

In particular, we study the well-posedness and the regularity of solutions. We also prove the dissipativity of the corresponding solution operators, as well as the existence of the global attractor. We finally give numerical simulations which show the effects of the higher-order terms on the anisotropy.

2 Setting of the problem

We consider the following initial and boundary value problem, for $k \in \mathbb{N}$, $k \geq 2$ (the case $k = 1$ can be treated as in [35]):

$$\frac{\partial u}{\partial t} - \Delta \sum_{i=1}^k (-1)^i \sum_{|\alpha|=i} a_\alpha \mathcal{D}^{2\alpha} u - \Delta f(u) + g(x, u) = 0, \quad (2.1)$$

$$\mathcal{D}^\alpha u = 0 \quad \text{on } \Gamma, \quad |\alpha| \leq k, \quad (2.2)$$

$$u|_{t=0} = u_0. \quad (2.3)$$

We assume that

$$a_\alpha > 0, \quad |\alpha| = k, \quad (2.4)$$

and we introduce the elliptic operator A_k defined by

$$\langle A_k v, w \rangle_{H^{-k}(\Omega), H_0^k(\Omega)} = \sum_{|\alpha|=k} a_\alpha ((\mathcal{D}^\alpha v, \mathcal{D}^\alpha w)), \quad (2.5)$$

where $H^{-k}(\Omega)$ is the topological dual of $H_0^k(\Omega)$. Furthermore, $((\cdot, \cdot))$ denotes the usual L^2 -scalar product, with associated norm $\|\cdot\|$. More generally, we denote by $\|\cdot\|_X$ the norm on the Banach space X ; we also set $\|\cdot\|_{-1} = \|(-\Delta)^{-\frac{1}{2}} \cdot\|$, where $(-\Delta)^{-1}$ denotes the inverse minus Laplace operator associated with Dirichlet boundary conditions. We can note that

$$(v, w) \in H_0^k(\Omega)^2 \mapsto \sum_{|\alpha|=k} a_\alpha ((\mathcal{D}^\alpha v, \mathcal{D}^\alpha w))$$

is bilinear, symmetric, continuous and coercive, so that

$$A_k : H_0^k(\Omega) \rightarrow H^{-k}(\Omega)$$

is indeed well defined. It then follows from elliptic regularity results for linear elliptic operators of order $2k$ (see [1–3]) that A_k is a strictly positive, selfadjoint and unbounded linear operator with compact inverse, with domain

$$D(A_k) = H^{2k}(\Omega) \cap H_0^k(\Omega),$$

where, for $v \in D(A_k)$,

$$A_k v = (-1)^k \sum_{|\alpha|=k} a_\alpha \mathcal{D}^{2\alpha} v.$$

We further note that $D(A_k^{\frac{1}{2}}) = H_0^k(\Omega)$ and, for $(v, w) \in D(A_k^{\frac{1}{2}})^2$,

$$\left(\left(A_k^{\frac{1}{2}} v, A_k^{\frac{1}{2}} w \right) \right) = \sum_{|\alpha|=k} a_\alpha ((\mathcal{D}^\alpha v, \mathcal{D}^\alpha w)).$$

We finally note that (see, e.g., [43]) $\|A_k \cdot\|$ (resp., $\|A_k^{\frac{1}{2}} \cdot\|$) is equivalent to the usual H^{2k} -norm (resp., H^k -norm) on $D(A_k)$ (resp., $D(A_k^{\frac{1}{2}})$).

Similarly, we can define the linear operator $\bar{A}_k = -\Delta A_k$,

$$\bar{A}_k : H_0^{k+1}(\Omega) \rightarrow H^{-k-1}(\Omega)$$

which is a strictly positive, selfadjoint and unbounded linear operator with compact inverse, with domain

$$D(\bar{A}_k) = H^{2k+2}(\Omega) \cap H_0^{k+1}(\Omega),$$

where, for $v \in D(\bar{A}_k)$,

$$\bar{A}_k v = (-1)^{k+1} \Delta \sum_{|\alpha|=k} a_\alpha \mathcal{D}^{2\alpha} v.$$

Furthermore, $D(\bar{A}_k^{\frac{1}{2}}) = H_0^{k+1}(\Omega)$ and, for $(v, w) \in D(\bar{A}_k^{\frac{1}{2}})^2$,

$$\left(\left(\bar{A}_k^{\frac{1}{2}} v, \bar{A}_k^{\frac{1}{2}} w \right) \right) = \sum_{|\alpha|=k} a_\alpha ((\nabla \mathcal{D}^\alpha v, \nabla \mathcal{D}^\alpha w)).$$

Besides, $\|\bar{A}_k \cdot\|$ (resp., $\|\bar{A}_k^{\frac{1}{2}} \cdot\|$) is equivalent to the usual H^{2k+2} -norm (resp., H^{k+1} -norm) on $D(\bar{A}_k)$ (resp., $D(\bar{A}_k^{\frac{1}{2}})$).

We finally consider the operator $\tilde{A}_k = (-\Delta)^{-1} A_k$, where

$$\tilde{A}_k : H_0^{k-1}(\Omega) \rightarrow H^{-k+1}(\Omega);$$

note that, as $-\Delta$ and A_k commute, then the same holds for $(-\Delta)^{-1}$ and A_k , so that $\tilde{A}_k = A_k(-\Delta)^{-1}$.

We have the following lemma.

Lemma 2.1. *The operator \tilde{A}_k is a strictly positive, selfadjoint and unbounded linear operator with compact inverse, with domain*

$$D(\tilde{A}_k) = H^{2k-2}(\Omega) \cap H_0^{k-1}(\Omega),$$

where, for $v \in D(\tilde{A}_k)$,

$$\tilde{A}_k v = (-1)^k \sum_{|\alpha|=k} a_\alpha \mathcal{D}^{2\alpha} (-\Delta)^{-1} v.$$

Furthermore, $D(\tilde{A}_k^{\frac{1}{2}}) = H_0^{k-1}(\Omega)$ and, for $(v, w) \in D(\tilde{A}_k^{\frac{1}{2}})^2$,

$$\left(\left(\tilde{A}_k^{\frac{1}{2}} v, \tilde{A}_k^{\frac{1}{2}} w \right) \right) = \sum_{|\alpha|=k} a_\alpha \left(\left(\mathcal{D}^\alpha (-\Delta)^{-\frac{1}{2}} v, \mathcal{D}^\alpha (-\Delta)^{-\frac{1}{2}} w \right) \right).$$

Besides, $\|\tilde{A}_k \cdot\|$ (resp., $\|\tilde{A}_k^{\frac{1}{2}} \cdot\|$) is equivalent to the usual H^{2k-2} -norm (resp., H^{k-1} -norm) on $D(\tilde{A}_k)$ (resp., $D(\tilde{A}_k^{\frac{1}{2}})$).

Proof. We first note that \tilde{A}_k clearly is linear and unbounded. Then, since $(-\Delta)^{-1}$ and A_k commute, it easily follows that \tilde{A}_k is selfadjoint.

Next, the domain of \tilde{A}_k is defined by

$$D(\tilde{A}_k) = \left\{ v \in H_0^{k-1}(\Omega), \tilde{A}_k v \in L^2(\Omega) \right\}.$$

Noting that $\tilde{A}_k v = f$, $f \in L^2(\Omega)$, $v \in D(\tilde{A}_k)$, is equivalent to $A_k v = -\Delta f$, where $-\Delta f \in H^2(\Omega)'$, it follows from the elliptic regularity results of [1], [2] and [3] that $v \in H^{2k-2}(\Omega)$, so that $D(\tilde{A}_k) = H^{2k-2}(\Omega) \cap H_0^{k-1}(\Omega)$.

Noting then that \tilde{A}_k^{-1} maps $L^2(\Omega)$ onto $H^{2k-2}(\Omega)$ and recalling that $k \geq 2$, we deduce that \tilde{A}_k has compact inverse.

We now note that, considering the spectral properties of $-\Delta$ and A_k (see, e.g., [43]) and recalling that these two operators commute, $-\Delta$ and A_k have a spectral basis formed of common eigenvectors. This yields that, $\forall s_1, s_2 \in \mathbb{R}$, $(-\Delta)^{s_1}$ and $A_k^{s_2}$ commute.

Having this, we see that $\tilde{A}_k^{\frac{1}{2}} = (-\Delta)^{-\frac{1}{2}} A_k^{\frac{1}{2}}$, so that $D(\tilde{A}_k^{\frac{1}{2}}) = H_0^{k-1}(\Omega)$, and, for $(v, w) \in D(\tilde{A}_k^{\frac{1}{2}})^2$,

$$\left(\left(\tilde{A}_k^{\frac{1}{2}} v, \tilde{A}_k^{\frac{1}{2}} w \right) \right) = \sum_{|\alpha|=k} a_\alpha \left(\left(\mathcal{D}^\alpha (-\Delta)^{-\frac{1}{2}} v, \mathcal{D}^\alpha (-\Delta)^{-\frac{1}{2}} w \right) \right).$$

Finally, as far as the equivalences of norms are concerned, we can note that, for instance, the norm $\|\tilde{A}_k^{\frac{1}{2}} \cdot\|$ is equivalent to the norm $\|(-\Delta)^{-\frac{1}{2}} \cdot\|_{H^k(\Omega)}$ and, thus, to the norm $\|(-\Delta)^{\frac{k-1}{2}} \cdot\|$. \square

Having this, we rewrite (2.1) as

$$\frac{\partial u}{\partial t} - \Delta A_k u - \Delta B_k u - \Delta f(u) + g(x, u) = 0, \quad (2.6)$$

where

$$B_k v = \sum_{i=1}^{k-1} (-1)^i \sum_{|\alpha|=i} a_\alpha \mathcal{D}^{2\alpha} v.$$

As far as the nonlinear term f is concerned, we assume that

$$f \in \mathcal{C}^2(\mathbb{R}), \quad f(0) = 0, \quad (2.7)$$

$$f' \geq -c_0, \quad c_0 \geq 0, \quad (2.8)$$

$$f(s)s \geq c_1 F(s) - c_2 \geq -c_3, \quad c_1 > 0, \quad c_2, c_3 \geq 0, \quad s \in \mathbb{R}, \quad (2.9)$$

$$F(s) \geq c_4 s^4 - c_5, \quad c_4 > 0, \quad c_5 \geq 0, \quad s \in \mathbb{R}, \quad (2.10)$$

where $F(s) = \int_0^s f(\xi) d\xi$. In particular, the usual cubic nonlinear term $f(s) = s^3 - s$ satisfies these assumptions.

Furthermore, as far as the function g is concerned, we assume that

$$g(\cdot, s) \text{ is measurable, } \forall s \in \mathbb{R}, \quad g(x, \cdot) \text{ is of class } \mathcal{C}^1, \text{ a.e. } x \in \Omega, \quad (2.11)$$

$$\frac{\partial g}{\partial s}(\cdot, s) \text{ is measurable, } \quad \forall s \in \mathbb{R};$$

$$|g(x, s)| \leq h(s), \quad \text{a.e. } x \in \Omega, \quad s \in \mathbb{R}, \quad (2.12)$$

where $h \geq 0$ is continuous and satisfies

$$\|h(v)\| \|v\| \leq \varepsilon \int_{\Omega} F(v) dx + c_{\varepsilon}, \quad \forall \varepsilon > 0, \quad (2.13)$$

$\forall v \in L^2(\Omega)$ such that $\int_{\Omega} F(v) dx < +\infty$, and

$$|h(s)|^2 \leq c_6 F(s) + c_7, \quad c_6, c_7 \geq 0, \quad s \in \mathbb{R}; \quad (2.14)$$

$$\left| \frac{\partial g}{\partial s}(x, s) \right| \leq l(s), \quad \text{a.e. } x \in \Omega, \quad s \in \mathbb{R}, \quad (2.15)$$

where $l \geq 0$ is continuous.

Example 2.2. We assume that $f(s) = s^3 - s$. Assumptions (2.11)–(2.15) are satisfied in the following cases.

- (i) Cahn–Hilliard–Oono equation (see [34], [40] and [46]). In that case,

$$g(x, s) = g(s) = \beta s, \quad \beta > 0.$$

This function was proposed in [40] in order to account for long-ranged (i.e., nonlocal) interactions, but also to simplify numerical simulations.

- (ii) Proliferation term. In that case,

$$g(x, s) = g(s) = \beta s(s - 1), \quad \beta > 0.$$

This function was proposed in [26] in view of biological applications and, more precisely, to model wound healing and tumor growth (in one space dimension) and the clustering of brain tumor cells (in two space dimensions); see also [4] for other quadratic functions.

(iii) Fidelity term. In that case,

$$g(x, s) = \lambda_0 \chi_{\Omega \setminus D}(x)(s - \varphi(x)), \quad \lambda_0 > 0, \quad D \subset \Omega, \quad \varphi \in L^2(\Omega),$$

where χ denotes the indicator function. This function was proposed in [5] and [6] in view of applications to image inpainting. Here, φ is a given (damaged) image and D is the inpainting (i.e., damaged) region. Furthermore, the fidelity term $g(x, u)$ is added in order to keep the solution close to the image outside the inpainting region. The idea in this model is to solve the equation up to steady state to obtain an inpainted (i.e., restored) version $u(x)$ of $\varphi(x)$.

Throughout the paper, the same letters c , c' and c'' denote (generally positive) constants which may vary from line to line. Similarly, the same letters Q and Q' denote (positive) monotone increasing and continuous (with respect to each argument) functions which may vary from line to line.

3 A priori estimates

Proposition 3.1. *Any sufficiently regular solution to (2.1)–(2.3) satisfies the following estimates:*

$$\|u(t)\|_{H^k(\Omega)}^2 \leq ce^{-c't} \left(\|u_0\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u_0) dx \right) + c'', \quad c' > 0, \quad t \geq 0, \quad (3.1)$$

$$\int_t^{t+r} \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 ds \leq ce^{-c't} \left(\|u_0\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u_0) dx \right) + c'', \quad (3.2)$$

$c' > 0, \quad t \geq 0, \quad r > 0$ given,

and

$$\|u(t)\|_{H^{2k}(\Omega)} \leq Q(e^{-ct} Q'(\|u_0\|_{H^k(\Omega)}) + c'), \quad c > 0, \quad t \geq 1, \quad (3.3)$$

where the continuous and monotone increasing function Q is of the form $Q(s) = cse^{c's}$.

Proof. The estimates below will be formal, but they can easily be justified within, e.g., a standard Galerkin scheme.

We multiply (2.6) by $(-\Delta)^{-1} \frac{\partial u}{\partial t}$ and integrate over Ω and by parts. This gives

$$\frac{d}{dt} \left(\left\| A_k^{\frac{1}{2}} u \right\|^2 + B_k^{\frac{1}{2}}[u] + 2 \int_{\Omega} F(u) dx \right) + 2 \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 = - \left(\left(g(\cdot, u), (-\Delta)^{-1} \frac{\partial u}{\partial t} \right) \right),$$

where

$$B_k^{\frac{1}{2}}[u] = \sum_{i=1}^{k-1} \sum_{|\alpha|=i} a_{\alpha} \|\mathcal{D}^{\alpha} u\|^2$$

(note that $B_k^{\frac{1}{2}}[u]$ is not necessarily nonnegative). This yields, owing to (2.12) and (2.14),

$$\frac{d}{dt} \left(\left\| A_k^{\frac{1}{2}} u \right\|^2 + B_k^{\frac{1}{2}}[u] + 2 \int_{\Omega} F(u) dx \right) + \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 \leq c \int_{\Omega} F(u) dx + c'. \quad (3.4)$$

We can note that, owing to the interpolation inequality

$$\|v\|_{H^i(\Omega)} \leq c(i) \|v\|_{H^m(\Omega)}^{\frac{i}{m}} \|v\|^{1-\frac{i}{m}}, \quad (3.5)$$

$$v \in H^m(\Omega), \quad i \in \{1, \dots, m-1\}, \quad m \in \mathbb{N}, \quad m \geq 2,$$

there holds

$$|B_k^{\frac{1}{2}}[u]| \leq \frac{1}{2} \|A_k^{\frac{1}{2}} u\|^2 + c \|u\|^2.$$

This yields, employing (2.10),

$$\|A_k^{\frac{1}{2}} u\|^2 + B_k^{\frac{1}{2}}[u] + 2 \int_{\Omega} F(u) dx \geq \frac{1}{2} \|A_k^{\frac{1}{2}} u\|^2 + \int_{\Omega} F(u) dx + c \|u\|_{L^4(\Omega)}^4 - c' \|u\|^2 - c'',$$

whence

$$\|A_k^{\frac{1}{2}} u\|^2 + B_k^{\frac{1}{2}}[u] + 2 \int_{\Omega} F(u) dx \geq c \left(\|u\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u) dx \right) - c', \quad c > 0, \quad (3.6)$$

noting that, owing to Young's inequality,

$$\|u\|^2 \leq \varepsilon \|u\|_{L^4(\Omega)}^4 + c_{\varepsilon}, \quad \forall \varepsilon > 0. \quad (3.7)$$

We then multiply (2.6) by $(-\Delta)^{-1}u$ and have, owing to (2.9), (2.12), (2.13) and the interpolation inequality (3.5),

$$\frac{d}{dt} \|u\|_{-1}^2 + c \left(\|u\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u) dx \right) \leq c' \|u\|^2 + \varepsilon \int_{\Omega} F(u) dx + c''_{\varepsilon}, \quad \forall \varepsilon > 0,$$

hence, proceeding as above and employing, in particular, (2.10),

$$\frac{d}{dt} \|u\|_{-1}^2 + c \left(\|u\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u) dx \right) \leq c', \quad c > 0. \quad (3.8)$$

Summing δ_1 times (3.4) and (3.8), where $\delta_1 > 0$ is small enough, we obtain a differential inequality of the form

$$\frac{dE_1}{dt} + c \left(E_1 + \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 \right) \leq c', \quad c > 0, \quad (3.9)$$

where

$$E_1 = \delta_1 \left(\|A_k^{\frac{1}{2}} u\|^2 + B_k^{\frac{1}{2}}[u] + 2 \int_{\Omega} F(u) dx \right) + \|u\|_{-1}^2$$

satisfies, owing to (3.6),

$$E_1 \geq c \left(\|u\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u) dx \right) - c', \quad c > 0. \quad (3.10)$$

Note indeed that

$$\begin{aligned} E_1 &\leq c \|u\|_{H^k(\Omega)}^2 + 2 \int_{\Omega} F(u) dx \\ &\leq c \left(\|u\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u) dx \right) - c', \quad c > 0, \quad c' \geq 0. \end{aligned}$$

Estimates (3.1)–(3.2) then follow from (3.9)–(3.10) and Gronwall's lemma.

Multiplying next (2.6) by $\tilde{A}_k u$, we find, owing to (2.12) and the interpolation inequality (3.5),

$$\frac{d}{dt} \|\tilde{A}_k^{\frac{1}{2}} u\|^2 + c \|u\|_{H^{2k}(\Omega)}^2 \leq c (\|u\|^2 + \|f(u)\|^2 + \|h(u)\|^2). \quad (3.11)$$

It follows from the continuity of f , F and h , the continuous embedding $H^k(\Omega) \subset \mathcal{C}(\overline{\Omega})$ (recall that $k \geq 2$) and (3.1) that

$$\|u\|^2 + \|f(u)\|^2 + \|h(u)\|^2 \leq Q(\|u\|_{H^k(\Omega)}) \leq e^{-ct} Q'(\|u_0\|_{H^k(\Omega)}) + c', \quad c > 0, t \geq 0, \quad (3.12)$$

so that

$$\frac{d}{dt} \left\| \tilde{A}_k^{\frac{1}{2}} u \right\|^2 + c \|u\|_{H^{2k}(\Omega)}^2 \leq e^{-c't} Q(\|u_0\|_{H^k(\Omega)}) + c'', \quad c, c' > 0, t \geq 0. \quad (3.13)$$

Summing (3.9) and (3.13), we have a differential inequality of the form

$$\frac{dE_2}{dt} + c \left(E_2 + \|u\|_{H^{2k}(\Omega)}^2 + \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 \right) \leq e^{-c't} Q(\|u_0\|_{H^k(\Omega)}) + c'', \quad c, c' > 0, t \geq 0, \quad (3.14)$$

where

$$E_2 = E_1 + \|\tilde{A}_k^{\frac{1}{2}} u\|^2$$

satisfies

$$E_2 \geq c \left(\|u\|_{H^k(\Omega)}^2 + \int_{\Omega} F(u) dx \right) - c', \quad c > 0. \quad (3.15)$$

We now multiply (2.6) by $\frac{\partial u}{\partial t}$ and obtain, noting that f is of class \mathcal{C}^2 , so that

$$\|\Delta f(u)\| \leq Q(\|u\|_{H^k(\Omega)}),$$

and proceeding as above,

$$\frac{d}{dt} \left(\left\| \tilde{A}_k^{\frac{1}{2}} u \right\|^2 + \overline{B}_k^{\frac{1}{2}}[u] \right) + \left\| \frac{\partial u}{\partial t} \right\|^2 \leq e^{-c't} Q(\|u_0\|_{H^k(\Omega)}) + c'', \quad c, c' > 0, \quad (3.16)$$

where

$$\overline{B}_k^{\frac{1}{2}}[u] = \sum_{i=1}^{k-1} \sum_{|\alpha|=i} a_{\alpha} \|\nabla \mathcal{D}^{\alpha} u\|^2.$$

Summing finally (3.14) and (3.16), we find a differential inequality of the form

$$\frac{dE_3}{dt} + c \left(E_3 + \|u\|_{H^{2k}(\Omega)}^2 + \left\| \frac{\partial u}{\partial t} \right\|^2 \right) \leq e^{-c't} Q(\|u_0\|_{H^k(\Omega)}) + c'', \quad c, c' > 0, t \geq 0, \quad (3.17)$$

where

$$E_3 = E_2 + \left\| \tilde{A}_k^{\frac{1}{2}} u \right\|^2 + \overline{B}_k^{\frac{1}{2}}[u]$$

satisfies, proceeding as above,

$$E_3 \geq c \left(\|u\|_{H^{k+1}(\Omega)}^2 + \int_{\Omega} F(u) dx \right) - c', \quad c > 0. \quad (3.18)$$

In particular, it follows from (3.17)–(3.18) that

$$\|u(t)\|_{H^{k+1}(\Omega)} \leq e^{-ct} Q(\|u_0\|_{H^{k+1}(\Omega)}) + c', \quad c > 0, t \geq 0. \quad (3.19)$$

We then rewrite (2.6) as an elliptic equation, for $t > 0$ fixed,

$$A_k u = -(-\Delta)^{-1} \frac{\partial u}{\partial t} - B_k u - f(u) - (-\Delta)^{-1} g(x, u), \quad \mathcal{D}^{\alpha} u = 0 \text{ on } \Gamma, \quad |\alpha| \leq k-1. \quad (3.20)$$

Multiplying (3.20) by $A_k u$, we have, owing to (2.12) and the interpolation inequality (3.5),

$$\|A_k u\|^2 \leq c \left(\|u\|^2 + \|f(u)\|^2 + \|h(u)\|^2 + \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 \right), \quad (3.21)$$

hence, proceeding as above (employing, in particular, (3.12)),

$$\|u\|_{H^{2k}(\Omega)}^2 \leq c \left(e^{-c't} Q(\|u_0\|_{H^k(\Omega)}) + \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 \right) + c'', \quad c' > 0. \quad (3.22)$$

In a next step, we differentiate (2.6) with respect to time and obtain

$$\frac{\partial}{\partial t} \frac{\partial u}{\partial t} - \Delta A_k \frac{\partial u}{\partial t} - \Delta B_k \frac{\partial u}{\partial t} - \Delta \left(f'(u) \frac{\partial u}{\partial t} \right) + \frac{\partial g}{\partial s}(x, u) \frac{\partial u}{\partial t} = 0, \quad (3.23)$$

$$\mathcal{D}^\alpha \frac{\partial u}{\partial t} = 0 \quad \text{on } \Gamma, \quad |\alpha| \leq k. \quad (3.24)$$

We multiply (3.23) by $(-\Delta)^{-1} \frac{\partial u}{\partial t}$ and find, owing to (2.8), (2.15), the interpolation inequality (3.5) and the continuous embedding $H^2(\Omega) \subset L^\infty(\Omega)$,

$$\begin{aligned} \frac{d}{dt} \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 + c \left\| \frac{\partial u}{\partial t} \right\|_{H^k(\Omega)}^2 &\leq c' \left(\left\| \frac{\partial u}{\partial t} \right\|^2 + \|l(u)\| \left\| \frac{\partial u}{\partial t} \right\| \left\| (-\Delta)^{-1} \frac{\partial u}{\partial t} \right\|_{L^\infty(\Omega)} \right) \\ &\leq c' \left(\left\| \frac{\partial u}{\partial t} \right\|^2 + \|l(u)\| \left\| \frac{\partial u}{\partial t} \right\|^2 \right), \quad c > 0, \end{aligned}$$

which yields, employing the interpolation inequality

$$\|v\|^2 \leq c \|v\|_{-1} \|v\|_{H^1(\Omega)}, \quad v \in H_0^1(\Omega), \quad (3.25)$$

and proceeding as above (note that l is continuous), the differential inequality

$$\frac{d}{dt} \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2 + c \left\| \frac{\partial u}{\partial t} \right\|_{H^k(\Omega)}^2 \leq c' (e^{-c''t} Q(\|u_0\|_{H^k(\Omega)}) + 1) \left\| \frac{\partial u}{\partial t} \right\|_{-1}^2, \quad c, c'' > 0. \quad (3.26)$$

In particular, this yields, owing to (3.2) and employing the uniform Gronwall's lemma (see, e.g., [43]),

$$\left\| \frac{\partial u}{\partial t}(t) \right\|_{-1} \leq \frac{1}{r^{\frac{1}{2}}} Q(e^{-ct} Q'(\|u_0\|_{H^k(\Omega)}) + c'), \quad c > 0, \quad t \geq r, \quad r > 0 \text{ given}. \quad (3.27)$$

Finally, (3.3) follows from (3.22) and (3.27) (for $r = 1$). \square

Remark 3.2. If we assume that $u_0 \in H^{2k+1}(\Omega) \cap H_0^k(\Omega)$, we deduce from (3.22), (3.26) and Gronwall's lemma an H^{2k} -estimate on u on $[0, 1]$ which, combined with (3.3), gives an H^{2k} -estimate on u , for all times. This is however not satisfactory, in particular, in view of the study of attractors.

Remark 3.3. We assume that, for simplicity, $g(x, s) = g(s)$ and we further assume that f is of class \mathcal{C}^{k+1} and g is of class \mathcal{C}^{k-1} . Multiplying (2.6) by $\tilde{A}_k \frac{\partial u}{\partial t}$, we have

$$\frac{1}{2} \frac{d}{dt} (\|A_k u\|^2 + ((A_k u, B_k u))) + \left\| \tilde{A}_k^{\frac{1}{2}} \frac{\partial u}{\partial t} \right\|^2 = - \left(\left(\tilde{A}_k^{\frac{1}{2}} f(u), \tilde{A}_k^{\frac{1}{2}} \frac{\partial u}{\partial t} \right) \right) - \left(\left(\tilde{A}_k^{\frac{1}{2}} g(u), \tilde{A}_k^{\frac{1}{2}} \frac{\partial u}{\partial t} \right) \right),$$

which yields, noting that $\|\bar{A}_k^{\frac{1}{2}}f(u)\|^2 + \|\bar{A}_k^{\frac{1}{2}}g(u)\|^2 \leq Q(\|u\|_{H^{k+1}(\Omega)})$ and owing to (3.19),

$$\frac{d}{dt}(\|A_k u\|^2 + ((A_k u, B_k u))) \leq e^{-ct} Q(\|u_0\|_{H^{k+1}(\Omega)}) + c', \quad c > 0, t \geq 0. \quad (3.28)$$

Combining (3.28) with (3.17), it follows from (3.18) and the interpolation inequality (3.5) that

$$\|u(t)\|_{H^{2k}(\Omega)} \leq Q(\|u_0\|_{H^{2k}(\Omega)}), \quad t \in [0, 1],$$

so that, owing to (3.3),

$$\|u(t)\|_{H^{2k}(\Omega)} \leq Q(e^{-ct} Q'(\|u_0\|_{H^{2k}(\Omega)}) + c'), \quad c > 0, t \geq 0. \quad (3.29)$$

4 The dissipative semigroup

We first give the definition of a weak solution to (2.1)–(2.3).

Definition 4.1. We assume that $u_0 \in L^2(\Omega)$. A weak solution to (2.1)–(2.3) is a function u such that, for any given $T > 0$,

$$u \in \mathcal{C}([0, T]; L^2(\Omega)) \cap L^2(0, T; H_0^k(\Omega)),$$

$$u(0) = u_0 \quad \text{in } L^2(\Omega)$$

and

$$\begin{aligned} \frac{d}{dt}(((-\Delta)^{-1}u, v)) + \sum_{i=1}^k \sum_{|\alpha|=i} a_i((\mathcal{D}^\alpha u, \mathcal{D}^\alpha v)) + ((f(u), v)) \\ + (((-\Delta)^{-1}g(x, u), v)) = 0, \quad \forall v \in H_0^k(\Omega), \end{aligned}$$

in the sense of distributions.

We have the following theorem.

Theorem 4.2.

- (i) We assume that $u_0 \in H_0^k(\Omega)$. Then, (2.1)–(2.3) possesses a unique weak solution u such that, $\forall T > 0$,

$$u \in L^\infty(\mathbb{R}^+; H_0^k(\Omega)) \cap L^2(0, T; H^{2k}(\Omega) \cap H_0^k(\Omega))$$

and

$$\frac{\partial u}{\partial t} \in L^2(0, T; H^{-1}(\Omega)).$$

- (ii) If we further assume that $u_0 \in H^{k+1}(\Omega) \cap H_0^k(\Omega)$, then, $\forall T > 0$,

$$u \in L^\infty(\mathbb{R}^+; H^{k+1}(\Omega) \cap H_0^k(\Omega))$$

and

$$\frac{\partial u}{\partial t} \in L^2(0, T; L^2(\Omega)).$$

- (iii) If we further assume that f is of class \mathcal{C}^{k+1} , $g(x, s) = g(s)$, g is of class \mathcal{C}^{k-1} and $u_0 \in H^{2k}(\Omega) \cap H_0^k(\Omega)$, then

$$u \in L^\infty(\mathbb{R}^+; H^{2k}(\Omega) \cap H_0^k(\Omega)).$$

Proof. The proofs of existence and regularity in (i), (ii) and (iii) follow from the a priori estimates derived in the previous section and, e.g., a standard Galerkin scheme. Indeed, we can note that, since the operators $-\Delta$, A_k , \bar{A}_k and \tilde{A}_k are linear, selfadjoint and strictly positive operators with compact inverse which commute, they have a spectral basis formed of common eigenvectors. We then take this spectral basis as Galerkin basis, so that all the a priori estimates derived in the previous section are justified within the Galerkin scheme.

Let now u_1 and u_2 be two solutions to (2.1)–(2.2) with initial data $u_{0,1}$ and $u_{0,2}$, respectively. We set $u = u_1 - u_2$ and $u_0 = u_{0,1} - u_{0,2}$ and have

$$\frac{\partial u}{\partial t} - \Delta A_k u - \Delta B_k u - \Delta(f(u_1) - f(u_2)) + g(x, u_1) - g(x, u_2) = 0, \quad (4.1)$$

$$\mathcal{D}^\alpha u = 0 \quad \text{on } \Gamma, \quad |\alpha| \leq k, \quad (4.2)$$

$$u|_{t=0} = u_0. \quad (4.3)$$

Multiplying (4.1) by $(-\Delta)^{-1}u$, we obtain, owing to (2.8), (2.15), (3.1) and the interpolation inequalities (3.5) and (3.25),

$$\frac{d}{dt} \|u\|_{-1}^2 + c \|u\|_{H^k(\Omega)}^2 \leq Q \|u\|_{-1}^2, \quad c > 0, \quad (4.4)$$

where

$$Q = Q(\|u_{0,1}\|_{H^k(\Omega)}, \|u_{0,2}\|_{H^k(\Omega)}).$$

Here, we have used the fact that, owing to (2.15) and (3.1),

$$\begin{aligned} \|g(x, u_1) - g(x, u_2)\| &\leq Q(\|u_1\|_{H^k(\Omega)}, \|u_2\|_{H^k(\Omega)}) \|u\| \\ &\leq Q(\|u_{0,1}\|_{H^k(\Omega)}, \|u_{0,2}\|_{H^k(\Omega)}) \|u\|. \end{aligned}$$

It follows from (4.4) and Gronwall's lemma that

$$\|u(t)\|_{-1}^2 \leq e^{Qt} \|u_0\|_{-1}^2, \quad t \geq 0, \quad (4.5)$$

hence the uniqueness, as well as the continuous dependence with respect to the initial data in the H^{-1} -norm. \square

It follows from Theorem 4.2 that we can define the family of solving operators

$$S(t) : \Phi \rightarrow \Phi, \quad u_0 \mapsto u(t), \quad t \geq 0,$$

where $\Phi = H_0^k(\Omega)$. This family of solving operators forms a semigroup which is continuous with respect to the H^{-1} -topology. Finally, it follows from (3.1) that we have the following theorem.

Theorem 4.3. *The semigroup $S(t)$ is dissipative in Φ , in the sense that it possesses a bounded absorbing set $\mathcal{B}_0 \subset \Phi$ (i.e., $\forall B \subset \Phi$ bounded, $\exists t_0 = t_0(B) \geq 0$ such that $t \geq t_0 \implies S(t)B \subset \mathcal{B}_0$).*

Remark 4.4.

- (i) Actually, it follows from (3.3) that we have a bounded absorbing set \mathcal{B}_1 which is compact in Φ and bounded in $H^{2k}(\Omega)$. This yields the existence of the global attractor \mathcal{A} which is compact in Φ and bounded in $H^{2k}(\Omega)$.
- (ii) We recall that the global attractor \mathcal{A} is the smallest (for the inclusion) compact set of the phase space which is invariant by the flow (i.e., $S(t)\mathcal{A} = \mathcal{A}$, $\forall t \geq 0$) and attracts all bounded sets of initial data as time goes to infinity; it thus appears as a suitable object in view of the study of the asymptotic behavior of the system. We refer the reader to, e.g., [37] and [43] for more details and discussions on this.

- (iii) We can also prove, based on standard arguments (see, e.g., [37] and [43]) that \mathcal{A} has finite dimension, in the sense of covering dimensions such as the Hausdorff and the fractal dimensions. The finite-dimensionality means, very roughly speaking, that, even though the initial phase space has infinite dimension, the reduced dynamics can be described by a finite number of parameters (we refer the interested reader to, e.g., [37] and [43] for discussions on this subject).

Remark 4.5. In the numerical simulations given in the next section below, the equations will be endowed with periodic boundary conditions. From a mathematical point of view, these boundary conditions are much more delicate to handle, since we have to estimate the spatial average of the order parameter $\langle u \rangle = \frac{1}{\text{Vol}(\Omega)} \int_{\Omega} u \, dx$ (see [12], [16] and [21]). When $g \equiv 0$, this is straightforward, since we have the conservation of mass, namely,

$$\langle u(t) \rangle = \langle u_0 \rangle, \quad \forall t \geq 0.$$

However, when g does not vanish, we are not able to estimate this quantity in general.

5 Numerical simulations

We give in this section several numerical simulations in order to illustrate the effects of the higher-order terms on the anisotropy. The computations presented below are performed with the software FreeFem++ (see [24]), for $k = 2$. We also take Ω bi-dimensional and rectangular. Finally, the system is associated with periodic boundary conditions.

The problem can be written as, for $k = 2$,

$$\begin{cases} \frac{\partial u}{\partial t} + \Delta w + \frac{1}{\varepsilon} g(x, u) = 0, \\ w + a_{20} \varepsilon \frac{\partial^4 u}{\partial x^4} + a_{02} \varepsilon \frac{\partial^4 u}{\partial y^4} + a_{11} \varepsilon \frac{\partial^4 u}{\partial x^2 \partial y^2} - a_{10} \varepsilon \frac{\partial^2 u}{\partial x^2} - a_{01} \varepsilon \frac{\partial^2 u}{\partial y^2} + \frac{1}{\varepsilon} f(u) = 0, \\ u, w \text{ are } \Omega\text{-periodic}, \\ u(0, x, y) = u_0(x, y), \end{cases}$$

where $\varepsilon > 0$ is introduced to take into account the diffuse interface thickness. Setting

$$\frac{\partial^2 u}{\partial x^2} = p, \quad \frac{\partial^2 u}{\partial y^2} = q, \quad \frac{\partial^4 u}{\partial x^2 \partial y^2} = \frac{1}{2} \frac{\partial^2 p}{\partial y^2} + \frac{1}{2} \frac{\partial^2 q}{\partial x^2},$$

we have the variational formulation: find $(u, w, p, q) \in H_{\text{per}}^1(\Omega)^4$ such that

$$\begin{cases} ((\frac{\partial u}{\partial t}, v_1)) - ((\nabla w, \nabla v_1)) + \frac{1}{\varepsilon} ((g(x, u), v_1)) = 0, \\ ((w, v_2)) - a_{20} \varepsilon ((\frac{\partial p}{\partial x}, \frac{\partial v_2}{\partial x})) - a_{02} \varepsilon ((\frac{\partial q}{\partial y}, \frac{\partial v_2}{\partial y})) - \frac{a_{11} \varepsilon}{2} ((\frac{\partial p}{\partial y}, \frac{\partial v_2}{\partial y})) \\ \quad - \frac{a_{11} \varepsilon}{2} ((\frac{\partial q}{\partial x}, \frac{\partial v_2}{\partial x})) - a_{10} \varepsilon ((p, v_2)) - a_{01} \varepsilon ((q, v_2)) + \frac{1}{\varepsilon} ((f(u), v_2)) = 0, \\ ((p, v_3)) + ((\frac{\partial u}{\partial x}, \frac{\partial v_3}{\partial x})) = 0, \\ ((q, v_4)) + ((\frac{\partial u}{\partial y}, \frac{\partial v_4}{\partial y})) = 0, \end{cases}$$

where the test functions v_1, v_2, v_3, v_4 all belong to $H_{\text{per}}^1(\Omega)$.

The mesh is obtained by dividing Ω into 149^2 rectangles, each rectangle being divided along the same diagonal into two triangles. The computations in Fig. 5.2, 5.3, 5.4 are based on a P_1 finite element method for the space discretization, while we used a P_2 finite element

method for Fig. 5.5, 5.6, 5.7. The time discretization uses a semi-implicit Euler scheme (implicit for the linear terms and explicit for the nonlinear ones).

We give numerical results concerning a higher-order Cahn–Hilliard–Oono equation (Fig. 5.2), a higher order phase-field crystal equation (Fig. 5.3, 5.4; see also [25]) and a higher-order Cahn–Hilliard equation with a mass source for tumor growth (Fig. 5.5, 5.6, 5.7; see also [4]). These results show that the anisotropy is strongly influenced by the choice of the coefficients in the higher-order terms. In particular, we can clearly see the anisotropy in the x , y and cross-directions. For instance, Fig. 5.5, Column 1, corresponds to a tumor growth simulated with the classical Cahn–Hilliard model (analogous simulations were performed in [4]). With very small coefficients for the sixth-order terms, the tumor evolves similarly, although the x , y and cross-directions are clearly noticeable (see Fig. 5.6). With larger coefficients, the tumor spreads and evolves faster; the anisotropy directions also become obvious (see Fig. 5.5, Column 2, for an isotropic situation and Fig. 5.7 for anisotropic ones).

(i) Cahn–Hilliard–Oono equation. (See Fig. 5.2.)

$$\begin{cases} f(u) = u^3 - u, & g(x, u) = 0.5u, & \varepsilon = 0.05, \\ u_0^{(1)} \text{ randomly distributed between } -1 \text{ and } 1, \\ \Omega = [0, 1] \times [0, 1], & \text{step size } \Delta t = 5 \times 10^{-8}, \\ \text{coefficients } a_{ij} \text{ in Table 5.1.} \end{cases}$$

(ii) Phase-field crystal equation. (See Fig. 5.3.)

$$\begin{cases} f(u) = u^3 + (1 - 0.025)u, & g(x, u) = 2u, & \varepsilon = 1, \\ u_0^{(2)} \text{ randomly distributed between } -0.2 \text{ and } 0.3, \\ \Omega = [-10, 10] \times [-10, 10], & \Delta t = 10^{-4}, \\ \text{coefficients } a_{ij} \text{ in Table 5.2.} \end{cases}$$

(iii) Phase-field crystal equation. (See Fig. 5.4.)

$$\begin{cases} f(u) = u^3 + (1 - 0.025)u, & g(x, u) = 2u, & \varepsilon = 1, \\ u_0^{(3)} = 0.07 - 0.02 \cos \frac{2\pi(x-12)}{32} \sin \frac{2\pi(y-1)}{32} \\ \quad + 0.02 \cos^2 \frac{\pi(x+10)}{32} \cos^2 \frac{\pi(y+3)}{32} \\ \quad - 0.01 \sin^2 \frac{4\pi x}{32} \sin^2 \frac{4\pi(y-6)}{32}, \\ \Omega = [0, 32] \times [0, 32], & \Delta t = 10^{-3}, \\ \text{coefficients } a_{ij} \text{ in Table 5.3.} \end{cases}$$

(iv) Tumor proliferation term. (See Fig. 5.5, 5.6, 5.7.)

$$\begin{cases} f(u) = u^3 - u, & \Omega = [-0.7, 1.7] \times [-1.7, 0.7], & \Delta t = 10^{-6} \\ g(x, u) = 46(u + 1) - 280(u - 1)^2(u + 1)^2, & \varepsilon = 0.0125, \\ u_0^{(4)} = -\tanh\left(\frac{1}{\sqrt{2\varepsilon}}\left(\sqrt{2(x - 0.5)^2 + 0.25(y + 0.5)^2} - 0.1\right)\right) \in [-1, 1], \\ \text{coefficients } a_{ij} \text{ in Tables 5.4, 5.5, 5.6.} \end{cases}$$

The initial conditions $u_0^{(3)}$ and $u_0^{(4)}$ are shown in Fig. 5.1.




Figure 5.1: Initial conditions $u_0^{(3)}$ and $u_0^{(4)}$.




Figure 5.2: Cahn–Hilliard–Oono. Initial condition $u_0^{(1)}$, $f = u^3 - u$, $g = 0.5u$, $\varepsilon = 0.05$, $\Delta t = 5 \times 10^{-8}$.

Table 5.1: Coefficients a_{ij} for Fig. 5.2.

column	a_{20}	a_{11}	a_{02}	a_{10}	a_{01}	Remark
1	0	0	0	1	1	Cahn–Hilliard–Oono
2	1e-2	1e-4	1e-4	1e-4	1e-4	x-direction
3	1e-4	1e-2	1e-4	1e-4	1e-4	cross-direction
4	1e-4	1e-4	1e-2	1e-4	1e-4	y-direction




Figure 5.3: Phase-field crystal. Initial condition $u_0^{(2)}$, $f = u^3 + (1 - 0.025)u$, $g = 2u$, $\varepsilon = 1$, $\Delta t = 10^{-4}$.

Table 5.2: Coefficients a_{ij} for Fig. 5.3.

Column	a_{20}	a_{11}	a_{02}	a_{10}	a_{01}	Remark
1	1	1	1	-2	-2	Phase-field crystal
2	1	0.1	0.1	-2	-2	x-direction
3	0.1	1	0.1	-2	-2	cross-direction
4	0.1	0.1	1	-2	-2	y-direction




Figure 5.4: Phase-field crystal. Initial condition $u_0^{(3)}$, $f = u^3 + (1 - 0.025)u$, $g = 2u$, $\varepsilon = 1$, $\Delta t = 10^{-3}$.

Table 5.3: Coefficients a_{ij} for Fig. 5.4.

Column	a_{20}	a_{11}	a_{02}	a_{10}	a_{01}	Remark
1	1	0.5	0.5	-2	-2	x-direction
2	0.5	1	0.5	-2	-2	cross-direction
3	0.5	0.5	1	-2	-2	y-direction




Figure 5.5: Tumor growth. Initial condition $u_0^{(4)}$, $f = u^3 - u$, $g = 46(u + 1) - 280(u - 1)^2(u + 1)^2$, $\varepsilon = 0.0125$, $\Delta t = 10^{-6}$.

Table 5.4: Coefficients a_{ij} for Fig. 5.5.

Column	a_{20}	a_{11}	a_{02}	a_{10}	a_{01}	Remark
1	0	0	0	1	1	Cahn–Hilliard isotropy
2	5e-5	5e-5	5e-5	1	1	




Figure 5.6: Tumor growth. Initial condition $u_0^{(4)}$, $f = u^3 - u$, $g = 46(u + 1) - 280(u - 1)^2(u + 1)^2$, $\varepsilon = 0.0125$, $\Delta t = 10^{-6}$.

Table 5.5: Coefficients a_{ij} for Fig. 5.6.

Column	a_{20}	a_{11}	a_{02}	a_{10}	a_{01}	Remark
1	1.8e-5	5e-6	5e-6	1	1	x-direction
2	5e-6	1.8e-5	5e-6	1	1	cross-direction
3	5e-6	5e-6	1.8e-5	1	1	y-direction




Figure 5.7: Tumor growth. Initial condition $u_0^{(4)}$, $f = u^3 - u$, $g = 46(u + 1) - 280(u - 1)^2(u + 1)^2$, $\varepsilon = 0.0125$, $\Delta t = 10^{-6}$.

Table 5.6: Coefficients a_{ij} for Fig. 5.7.

Column	a_{20}	a_{11}	a_{02}	a_{10}	a_{01}	Remark
1	5e-4	5e-6	5e-6	1	1	x-direction
2	5e-6	5e-4	5e-6	1	1	cross-direction
3	5e-6	5e-6	5e-4	1	1	y-direction

Acknowledgments

The authors wish to thank the Ministère Français des Affaires Etrangères et du Développement International (MAEDI), the Ministère de l'Education Nationale, de L'Enseignement Supérieur et de la Recherche (MENESR) and the China Scholarship Council (CSC) for their support within the PHC Program Cai Yuanpei 2015 "Modèles fortement anisotropes en séparation de phase".

References

- [1] S. AGMON, *Lectures on elliptic boundary value problems*, Van Nostrand Mathematical Studies, No. 2, Van Nostrand, New York, 1965. [MR178246](#)
- [2] S. AGMON, A. DOUGLIS, L. NIRENBERG, Estimates near the boundary for solutions of elliptic partial differential equations. I, *Commun. Pure Appl. Math.* **12**(1959), 623–727. [MR0125307](#); [url](#)
- [3] S. AGMON, A. DOUGLIS, L. NIRENBERG, Estimates near the boundary for solutions of elliptic partial differential equations. II, *Commun. Pure Appl. Math.* **17**(1964), 35–92. [MR0162050](#); [url](#)

- [4] A. C. ARISTOTELOUS, O. A. KARAKASHIAN, S. M. WISE, Adaptive, second-order in time, primitive-variable discontinuous Galerkin schemes for a Cahn–Hilliard equation with a mass source, *IMA J. Numer. Anal.* **35**(2015), 1167–1198. [MR3407258](#); [url](#)
- [5] A. BERTOZZI, S. ESEDOGLU, A. GILLETTE, Inpainting of binary images using the Cahn–Hilliard equation, *IEEE Trans. Imag. Proc.* **16**(2007), 285–291. [MR2460167](#); [url](#)
- [6] A. BERTOZZI, S. ESEDOGLU, A. GILLETTE, Analysis of a two-scale Cahn–Hilliard model for binary image inpainting, *Multiscale Model. Simul.* **6**(2007), 913–936. [MR2368973](#); [url](#)
- [7] G. CAGINALP, E. ESENTURK, Anisotropic phase field equations of arbitrary order, *Discrete Contin. Dyn. Systems Ser. S* **4**(2011), 311–350. [MR2746377](#); [url](#)
- [8] J. W. CAHN, On spinodal decomposition, *Acta Metall.* **9**(1961), 795–801. [url](#)
- [9] J. W. CAHN, J. E. HILLIARD, Free energy of a nonuniform system I. Interfacial free energy, *J. Chem. Phys.* **28**(1958), 258–267. [url](#)
- [10] V. CHALUPECKÝ, Numerical studies of Cahn–Hilliard equations and applications in image processing, in: *Proceedings of Czech–Japanese Seminar in Applied Mathematics 2004 (August 4–7, 2004)*, Czech Technical University in Prague. [url](#)
- [11] X. CHEN, G. CAGINALP, E. ESENTURK, Interface conditions for a phase field model with anisotropic and non-local interactions, *Arch. Rational Mech. Anal.* **202**(2011), 349–372. [MR2847529](#); [url](#)
- [12] L. CHERFILS, H. FAKIH, A. MIRANVILLE, Finite-dimensional attractors for the Bertozzi–Esedoglu–Gillette–Cahn–Hilliard equation in image inpainting, *Inv. Prob. Imag.* **9**(2015), 105–125. [MR3305888](#); [url](#)
- [13] L. CHERFILS, A. MIRANVILLE, S. PENG, Higher-order Allen–Cahn models with logarithmic nonlinear terms, in: V. A. Sadovnichiy, M. Z. Zgurovsky (eds.), *Advances in Dynamical Systems and Control*, Studies in Systems, Decision and Control, Vol. 69, Springer, 2016, pp. 247–263. [url](#)
- [14] L. CHERFILS, A. MIRANVILLE, S. PENG, Higher-order models in phase separation, *J. Appl. Anal. Comput.* **7**(2017), 39–56. [MR3528197](#); [url](#)
- [15] L. CHERFILS, A. MIRANVILLE, S. PENG, Higher-order anisotropic models in phase separation, submitted.
- [16] L. CHERFILS, A. MIRANVILLE, S. ZELIK, The Cahn–Hilliard equation with logarithmic potentials, *Milan J. Math.* **79**(2011), 561–596. [MR2862028](#); [url](#)
- [17] L. CHERFILS, A. MIRANVILLE, S. ZELIK, On a generalized Cahn–Hilliard equation with biological applications, *Discrete Cont. Dyn. Systems Ser. B* **19**(2014), 2013–2026. [MR3253242](#); [url](#)
- [18] D. COHEN, J. M. MURRAY, A generalized diffusion model for growth and dispersion in a population, *J. Math. Biol.* **12**(1981), 237–248. [url](#)
- [19] I. C. DOLCETTA, S. F. VITA, Area-preserving curve-shortening flows: from phase separation to image processing, *Interfaces Free Bound.* **4**(2002), 325–343. [MR1935642](#); [url](#)

- [20] C. M. ELLIOTT, The Cahn–Hilliard model for the kinetics of phase separation, in: J. F. Rodrigues (ed.), *Mathematical models for phase change problems*, International Series of Numerical Mathematics, Vol. 88, Birkhäuser, Basel, 1989. [MR1038064](#)
- [21] H. FAKIH, A Cahn–Hilliard equation with a proliferation term for biological and chemical applications, *Asympt. Anal.* **94**(2015), 71–104. [MR3385244](#); [url](#)
- [22] G. GIACOMIN, J. L. LEBOWITZ, Phase segregation dynamics in particle systems with long range interaction. I. Macroscopic limits, *J. Statist. Phys.* **87**(1997), 37–61. [MR1453735](#); [url](#)
- [23] G. GIACOMIN, J. L. LEBOWITZ, Phase segregation dynamics in particle systems with long range interaction. II. Interface motion, *SIAM J. Appl. Math.* **58**(1998), 1707–1729. [MR1638739](#); [url](#)
- [24] F. HECHT, New development in FreeFem++, *J. Numer. Math.* **20**(2012), 251–265. [MR3043640](#); [url](#)
- [25] Z. HU, S. M. WISE, C. WANG, J. S. LOWENGRUB, Stable and efficient finite-difference non linear multigrid schemes for the phase field crystal equation, *J. Comp. Phys.* **228**(2009), 5323–5339. [url](#)
- [26] E. KHAIN, L. M. SANDER, A generalized Cahn–Hilliard equation for biological applications, *Phys. Rev. E* **77**(2008), 051129. [url](#)
- [27] I. KLAPPER, J. DOCKERY, Role of cohesion in the material description of biofilms, *Phys. Rev. E* **74**(2006), 0319021. [url](#)
- [28] R. KOBAYASHI, Modelling and numerical simulations of dendritic crystal growth, *Phys. D* **63**(1993), 410–423. [url](#)
- [29] R. V. KOHN, F. OTTO, Upper bounds for coarsening rates, *Commun. Math. Phys.* **229**(2002), 375–395. [url](#)
- [30] J. S. LANGER, Theory of spinodal decomposition in alloys, *Ann. Phys.* **65**(1975), 53–86. [url](#)
- [31] Q.-X. LIU, A. DOELMAN, V. ROTTSCHÄFER, M. DE JAGER, P. M. J. HERMAN, M. RIETKERK, J. VAN DE KOPPEL, Phase separation explains a new class of self-organized spatial patterns in ecological systems, *Proc. Nation. Acad. Sci.* **110**(2013), No. 29, 11905–11910. [MR3104979](#); [url](#)
- [32] S. MAIER-PAAPE, T. WANNER, Spinodal decomposition for the Cahn–Hilliard equation in higher dimensions. I. Probability and wavelength estimate, *Commun. Math. Phys.* **195**(1998), 435–464. [MR1637817](#); [url](#)
- [33] S. MAIER-PAAPE, T. WANNER, Spinodal decomposition for the Cahn–Hilliard equation in higher dimensions: Nonlinear dynamics, *Arch. Ration. Mech. Anal.* **151**(2000), 187–219. [MR1753703](#); [url](#)
- [34] A. MIRANVILLE, Asymptotic behavior of the Cahn–Hilliard–Oono equation, *J. Appl. Anal. Comp.* **1**(2011), 523–536. [MR2889956](#); [url](#)
- [35] A. MIRANVILLE, Asymptotic behavior of a generalized Cahn–Hilliard equation with a proliferation term, *Appl. Anal.* **92**(2013), 1308–1321. [MR3197937](#); [url](#)

- [36] A. MIRANVILLE, A generalized Cahn–Hilliard equation with logarithmic potentials, in: V. A. Sadovnichiy, M. Z. Zgurovsky (eds.), *Continuous and Distributed Systems II*, Studies in Systems, Decision and Control, Vol. 30, Springer, 2015, pp. 137–148. [MR3381628](#); [url](#)
- [37] A. MIRANVILLE, S. ZELIK, Attractors for dissipative partial differential equations in bounded and unbounded domains, in: C. M. Dafermos, M. Pokorný (eds.), *Handbook of differential equations: evolutionary equations*, Vol. IV, Elsevier, Amsterdam, 2008, pp. 103–200. [MR2508165](#); [url](#)
- [38] A. NOVICK-COHEN, The Cahn–Hilliard equation: mathematical and modeling perspectives, *Adv. Math. Sci. Appl.* **8**(1998), 965–985. [MR1657208](#)
- [39] A. NOVICK-COHEN, The Cahn–Hilliard equation, in: C. M. Dafermos, M. Pokorný (eds.), *Handbook of differential equations: evolutionary equations*, Vol. IV, Elsevier, Amsterdam, 2008, pp. 201–228. [MR2508166](#); [url](#)
- [40] Y. OONO, S. PURI, Computationally efficient modeling of ordering of quenched phases, *Phys. Rev. Letters* **58**(1987), 836–839. [url](#)
- [41] A. ORON, S. H. DAVIS, S. G. BANKOFF, Long-scale evolution of thin liquid films, *Rev. Mod. Phys.* **69**(1997), 931–980. [url](#)
- [42] J. E. TAYLOR, II—mean curvature and weighted mean curvature, *Acta Metall. Mater.* **40**(1992), 1475–1495. [url](#)
- [43] R. TEMAM, *Infinite-dimensional dynamical systems in mechanics and physics*, Applied Mathematical Sciences, Vol. 68, Springer-Verlag, New York, 1997. [MR1441312](#)
- [44] U. THIELE, E. KNOBLOCH, Thin liquid films on a slightly inclined heated plate, *Phys. D* **190**(2004), 213–248. [MR2043346](#); [url](#)
- [45] S. TREMAINE, On the origin of irregular structure in Saturn’s rings, *Astron. J.* **125**(2003), 894–901. [url](#)
- [46] S. VILLAIN-GUILLOT, *Phases modulées et dynamique de Cahn–Hilliard*, Habilitation thesis, Université Bordeaux I, 2010.
- [47] A. A. WHEELER, G. B. MCFADDEN, On the notion of ζ -vector and stress tensor for a general class of anisotropic diffuse interface models, *Proc. R. Soc. London Ser. A* **453**(1997), 1611–1630. [url](#)